



Monkeypox

Monkeypox is a viral disease that normally starts with flu-like symptoms, including fever, chills, a headache, and achy muscles, with a rash usually then developing after 1 to 3 days. The rash typically progresses, beginning as flat spots that develop into bumps, which fill with fluid. Monkeypox is primarily spread through sustained physical contact such as skin-to-skin touch with someone who has an active rash. Respiratory spread is not the predominant worry.

Currently, there is no product on the market that has been tested against Monkeypox.

Emerging Viral Pathogen Claims:

Midlab® offers select **MAXIM**® products that qualify for emerging viral pathogen claims per the EPA's 'Guidance to Registrants: Process for Making Claims Against Emerging Viral Pathogens not on EPA-Registered Disinfectant Labels' when used in accordance with the appropriate use directions indicated in the chart below.

Product Name:	Reorder #	Follow the directions for use for the following viruses on the label:
Facility +	046200	Norovirus (Feline Calicivirus as the surrogate)Rhinovirus Type 37
Facility + RTU	046400	Norovirus (Feline Calicivirus as the surrogate)Rhinovirus Type 37
Citric Acid Disinfectant RTU	044600	Feline Calicivirus (ATCC VR-782) surrogate for Norovirus
Germicidal Cleaner	041000	Hepatitis A Virus
All Surface Bathroom Cleaner	038400	Hepatitis A Virus MAXIM
AFBC Acid Free Restroom Cleaner	036000	Hepatitis A Virus Hepatitis A Virus MAXIM MAXIM Secility * THE One-Step Disinfectant' Cleaner and Deodorant Contain hydrogen peroxide. For use with electrostatic sproyers. Facility * THE One-Step Disinfectant' Cleaner and Deodorant Contain hydrogen peroxide. For use with electrostatic sproyers. Facility * THE One-Step Disinfectant' Cleaner and Deodorant Contain hydrogen peroxide. For use with electrostatic sproyers. Facility * THE One-Step Disinfectant' Cleaner and Deodorant Contain hydrogen peroxide. For use with electrostatic sproyers. Facility * THE One-Step Disinfectant' Cleaner and Deodorant Contain hydrogen peroxide. For use with electrostatic sproyers. Facility * THE One-Step Disinfectant' Cleaner and Deodorant Contain hydrogen peroxide. For use with electrostatic sproyers. Facility * THE One-Step Disinfectant' Cleaner and Deodorant Contain hydrogen peroxide. For use with electrostatic sproyers. Facility * THE One-Step Disinfectant' Cleaner and Deodorant Contain hydrogen peroxide. For use with electrostatic sproyers. Facility * THE One-Step Disinfectant' Cleaner and Deodorant Contain hydrogen peroxide. For use with electrostatic sproyers. Facility * THE One-Step Disinfectant' Cleaner and Deodorant Contain hydrogen peroxide. For use with electrostatic sproyers. Facility * THE One-Step Disinfectant' Cleaner and Deodorant Contain hydrogen peroxide. For use with electrostatic sproyers. Facility * THE One-Step Disinfectant' Cleaner and Deodorant Contain hydrogen peroxide. For use with electrostatic sproyers. Facility * THE One-Step Disinfectant' Cleaner and Deodorant Contain hydrogen peroxide. For use with electrostatic sproyers. Facility * THE One-Step Disinfectant' Cleaner and Deodorant Contain hydrogen peroxide. For use with electrostatic sproyers. Facility * THE One-Step Disinfectant' Cleaner and Deodorant Contain hydrogen peroxide. For use with electrostatic sproyers. Facility * THE One-Step Disinfectant' Cleaner and Deodorant Facility * THE One-St
		Facility * TM RTU Dimfetant - Wirucidal Management and a real ma

midlab.com | 140 Private Brand Way, Athens, TN 37303 | 800.467.6294

Refer to the CDC website at https://www.cdc.gov/poxvirus/monkeypox/index.html for additional information.